

Simvastatin Induces Apoptosis of Fibroblast-Like Synoviocytes

Ira Litinsky*, Itshak Golan, Michael Yaron, Ilana Yaron, Dan Caspi and Ori Elkayam

Department of Rheumatology, Sourasky Medical Center and Sackler Faculty of Medicine, University of Tel Aviv, Tel Aviv, Israel

Abstract: *Background:* Statins (3-Hydroxy-3-methylglutaryl-CoA reductase inhibitors) exert favorable effects on lipoprotein metabolism, but appeared to possess anti-inflammatory properties among others, as suggested by their ability to inhibit collagen-induced arthritis in mice. Their activity in fibroblast-like synovial cells (FLS) has not yet been studied.

Objectives: To evaluate the effect of varying doses of simvastatin on apoptosis of FLS.

Methods: Synovial tissue, obtained during total knee replacement due to osteoarthritis, was cut into small pieces and cultured in Petri dishes with test materials, as previously described. FLS were incubated for 48 hours with 1 $\mu\text{mol/ml}$, 5 $\mu\text{mol/ml}$, 15 $\mu\text{mol/ml}$ and 50 $\mu\text{mol/ml}$ of simvastatin. Following incubation, apoptosis was analyzed by two-dimensional flow cytometry (FACS) using annexin V/PI staining according to the manufacturer's instructions.

Results: Different concentrations of simvastatin induced apoptosis of FLS. The level proportion of apoptotic cells of resting or activated with lipopolysaccharide (LPS; 3 $\mu\text{g/ml}$) FLS, not treated with simvastatin, was 21%. At 48 hours, the rate of apoptosis of activated fibroblasts, incubated with 1 $\mu\text{mol/ml}$, 5 $\mu\text{mol/ml}$, 15 and 50 $\mu\text{mol/ml}$ was 22%, 32%, 48% and 41% respectively. Synovial cell viability evaluated by tetrazolium salt XXT was unaffected by the simvastatin concentration used.

Conclusion: Varying concentrations of simvastatin induce apoptosis of activated fibroblast-like synoviocytes, suggesting another possible mechanism of anti-inflammatory effects of statins in inflammatory conditions.

Keywords: Apoptosis, inflammation, simvastatin, fibroblast-like synoviocytes.

INTRODUCTION

Statins are potent inhibitors of 3-hydroxy-3methylglutaryl coenzyme A (HMG-CoA) reductase, the rate-limiting enzyme of the cholesterol synthetic pathway, and they are widely used to treat hyperlipidemia. The beneficial effects of statins have been attributed primarily to their ability to inhibit cholesterol synthesis. They block the conversion of HMG-CoA to mevalonate at the start of the cholesterol biosynthetic pathway with a subsequent increase in LDL-receptor expression on the hepatocyte, resulting in a further reduction of plasma LDL-cholesterol level [1].

Although statins are considered to be the most effective drugs for treating hypercholesterolemia [2], the reduction in cardiovascular events observed in patients treated with statins is not fully explained by the reduction in LDL-cholesterol: this has been in part attributed to the ability of statins to modulate several mechanisms involved in the inflammatory process [3]. Better survival of heart transplant patients treated with statins has been attributed to their immunomodulatory effects [4] and reports of improvement of Rheumatoid Arthritis (RA) patients [5] lend support to the clinical anti-inflammatory role of statins.

Inflammatory processes are involved in atherosclerosis [3, 6]. A number of studies suggest that in addition to their

lipid-lowering activity, statins may influence atherosclerosis and cardiovascular diseases through the down regulation of the inflammatory response [7]. Statins has been shown to reduce plasma levels of TNF-alpha, sP selectin and IL-6 in long term survivors of heart transplantation and in patients with cholesterolemia, without known cardiovascular disease [8, 9].

Some *in vitro* experiments and studies in animal models have also found that these agents may down regulate inflammatory responses such as the production of pro-inflammatory cytokines, chemokines and adhesion molecules; these agents may also reduce the expression of inducible major histocompatibility complex (MHC) class II molecules by antigen presenting cells [10].

Rheumatoid arthritis (RA) is a chronic inflammatory joint disease characterized by synovial proliferation and abnormal immune responses, eventually causing erosion of the affected joint. Marked infiltration of lymphocytes, macrophages, and plasma cells is observed in the synovial tissue of RA [11]. Apoptosis of macrophages and fibroblast-like synoviocytes has been observed in rheumatoid synovium [12, 13]. It has been suggested that reduced apoptosis contributes to the synovial inflammatory process [12]. *In vitro* statins were shown to exert a cytotoxic effect on human T, B and myeloma cells by promoting their apoptosis [14] and also modifying apoptosis of smooth muscle and endothelial cells leading to altered vascular function and neovascularisation [15].

*Address correspondence to this author at the Department of Rheumatology, Sourasky Medical Center and Sackler Faculty of Medicine, University of Tel Aviv, 6 Weizmann Str, Tel Aviv 64239, Israel; Tel: +97236973668; Fax: +97236974577; E-mail: litinskyira@yahoo.com

We studied the effects of different doses of simvastatin on apoptosis of fibroblast-like synoviocytes (FLS).

MATERIALS AND METHODS

Specimen Selection and Culture Conditions

Synovial tissue was obtained during surgery from 3 patients undergoing total knee replacement because of osteoarthritis (OA). Synovial cells derived from small pieces (~2mm diameter) of human synovial tissue was first grown in T 15 culture flasks, by methods described by Castor *et al.* [16]. Then the cells identified as fibroblasts by the method of Castor [17] originating from the tissue samples were inoculated in T-15 culture flasks and incubated at 37°C. Cell were grown as monolayers in 6 or 7 day growth (between on trypsinization and another), then trypsinized and replated until there were enough cells for an experiment to be performed [17]. the culture medium consisted of RPMI 1640 supplemented by L-glutamine (2 mM) (Sigma, Israel), penicillin (100 U/ml), and streptomycin sulphate (100 U/ml) and 10% fetal calf serum (FCS, Bet-Haemek, Israel). Experiments were performed in 96 well plates containing $15\text{-}20\times 10^5$ cells/well in a serum free medium for 48 hours at 37°C (in 5% CO₂).

The LPS and simvastatin treatments were carried out with isolated cells in tissue culture cells and were incubated for 48 hours, using different concentrations of simvastatin: 1, 5, 15 and 50 µmol/ml. Stimulation of the cell culture (FLS) with lipopolysaccharide (LPS) was performed as previously described [18]. After collection of the cells, the incubation period of the cells with simvastatin lasted 48 hours.

The clinical research was conducted in accordance with guidelines for human experimentation specified by the Tel Aviv Sourasky Medical Center. The study was approved by the Ethics Committee of the Medical Center.

Apoptosis Assay

After 48 hours incubation in varying concentrations of simvastatin, fibroblast-like synovial cells were detached and placed in a serum free medium for 48 hours. Apoptosis was analyzed by two-dimensional flow cytometry, using annexin V and propidium iodide according to the manufacturer's instructions (Biotest). Flow cytometry was then performed using a Becton Dickinson FACSort flow cytometer (Becton Dickinson, San Jose, CA, USA). Events accumulating at the top and bottom right-hand quadrates simulated apoptotic cells.

Cell Viability and Toxicity Determination

Cell viability in the presence of statins was determined in human synovial cultures by tetrazolium salt XTT assay at the end of the experiment.

XTT is a tetrazolium derivative that measures cell viability based on the activity of mitochondria enzymes in live cells that reduce XTT and are inactivated shortly after cell death. XTT is readily reduced to a highly water-soluble orange colored. The amount of water-soluble product generated from XTT is proportional to the number of living cells in the sample and can be quantified by measuring absorbance at wavelength of 475 nm.

Statistical Analysis

Statistical analysis was evaluated by analysis of variance and by Student's t test. P values of less than 0.05 were considered statistically significant. Individual experiments were performed in triplicates. Results represent the findings from several separate experiments.

RESULTS

Apoptosis

Simvastatin induced apoptosis of human fibroblasts in a dose dependent manner. The proportion of apoptotic cells of resting or activated human fibroblasts lipopolysaccharide (LPS; 3µg/ml) not treated with simvastatin was 21% (Fig. 1). At 48 hours, the proportion of apoptosis of fibroblasts activated by LPS and incubated with 1, 5, 15 and 50 µmol/ml was 22%, 32%, 48% and 41% respectively (Fig. 2).

Cell Viability

Synovial cell viability evaluated by tetrazolium salt XTT was unaffected by the simvastatin concentration used in these experiments [19].

DISCUSSION

In this study, we have shown that simvastatin in doses within the therapeutic range causes a dose dependent induction of apoptosis of fibroblast-like synoviocytes activated with LPS. Despite the apoptotic effect of simvastatin, cell viability was not affected, suggesting that the level of apoptosis was limited. Our results are consistent with the data reported by Yokota *et al.* who have shown that high concentrations of simvastatin induce apoptosis of fibroblast-like-cells [20].

Inflammation of synovial tissue in rheumatoid arthritis is a complex sequence of events involving different cell lines and many soluble factors such as cytokines and chemokines.

A possible way to reduce the inflammatory burden may include an induction of apoptosis of cells involved in the inflammatory process. Several drugs, such as COX-2 inhibitors [21], methotrexate and corticosteroids [22] promote apoptosis of T cells. It has been recently shown that anti-tumor necrosis factor therapy with both etanercept and infliximab significantly increases apoptosis in the synovial monocyte/macrophage population [23].

Lovastatine and cerivastatin exert a cytostatic effect on mesothelioma [24], glioma [25], neoplastic thyroid [26], acute myeloid leukemia [27], multiple myeloma cells [28] and lung fibroblast [29] by directly promoting apoptosis. In myeloma cells, the apoptotic pathway engaged by statins is mediated by mitochondria activation, rather than by the "death receptors", and both caspases 3 and 9 appear to be directly involved and activated within the intrinsic pathway of apoptosis [30]. Statins also enhance the rate of apoptosis in neointimal smooth muscle cells [31]. Recently Nagashima *et al.* have shown that fluvastatin but not pravastatin, induced apoptosis in RA synoviocytes in a dose and time dependent manner [32]. Consistent with this report our study has demonstrated that simvastatin may also induce apoptosis of synovial cells.

In conclusion, our results demonstrate that simvastatin stimulates apoptosis of human synovial cells *in vitro*. We are aware of the limitations of this study which assessed apoptosis by Annexin PI staining without confirming the results using other methods. Further research is warranted to establish the *in vivo* relevance of these findings. Recently, increased cardiovascular morbidity and mortality have been observed in several inflammatory rheumatic diseases, including rheumatoid arthritis (RA) [35], encouraging aggressive treatment of the inflammatory process, as well as vigorous neutralization of traditional risk factors of atherosclerosis such as hypercholesterolemia. Our study further supports the use of statins in this subset of patients, suggesting an additional anti immunomodulatory effect.

REFERENCES

- [1] Aguilar-Salinas SA, Barrett H, Schonfeld G. Metabolic modes of action of statins in the hyperlipoproteinemias. *Atherosclerosis* 1998; 141: 203-7.
- [2] Downs JR, Clearfield M, Weis S, et al. Primary prevention of acute coronary events with lovastatin in men and women with average cholesterol levels: results of AFCAPS/TexCAPS. *JAMA* 1998; 279: 1615-22.
- [3] Case CC, Ballantyne CM. Statins and inflammatory markers. *Curr Atheroscler Rep* 2002; 4: 42-7.
- [4] Bilchick KC, Henrikson CA, Skojec D, Kasper EK, Blumenthal RS. Treatment of hyperlipidemia in cardiac transplant recipients. *Am Heart J* 2004; 148: 200-10.
- [5] McCarey DW, McInnes IB, Madhok R, et al. Trial of atorvastatin in rheumatoid arthritis (TARA): double-blind, randomized placebo-controlled trial. *Lancet* 2004; 363: 2015-21.
- [6] Ridker PM, Rifai N, Clearfield M, et al. Measurement of C-Reactive protein for the targeting of statin therapy in the primary prevention of acute coronary events. *N Engl J Med* 2001; 344: 1959-65.
- [7] Ridker PM, Rifai N, Lowenthal SP. Rapid reduction in C-reactive protein with cerivastatin among 785 patients with primary hypercholesterolemia. *Circulation* 2001; 103: 1191-3.
- [8] Torborn H, Andreassen AK, Ueland T, et al. Effect of pravastatin on plasma markers of inflammation and peripheral endothelial function in male heart transplant recipients. *Am J Cardio* 2001; 87: 815-18.
- [9] Rosenson RS, Tangney CC, Casey LC. Inhibition of proinflammatory cytokine production by pravastatin. *Lancet* 1999; 353: 983-84.
- [10] Jury EC, Ehrenstein MR. Statins: immunomodulators for autoimmune rheumatic disease? *Lupus* 2005; 14: 192-6.
- [11] Gay S, Gay RE, Koopman WJ. Molecular and cellular mechanisms of joint destruction in rheumatoid arthritis: two cellular mechanisms explain joint destruction. *Ann Rheum Dis* 1993; 52: S39-S47.
- [12] Firestein GS, Yeo M, Zvaifler NJ. Apoptosis in rheumatoid arthritis synovium. *J Clin Invest* 1995; 96: 1631-38.
- [13] Nagata S, Suda T. Fas and Fas ligand: lpr and gld mutations. *Immunol Today* 1995; 16: 39-43.
- [14] Cafforio P, Dammacco F, Gernone A, Silvestris F. Statins activate the mitochondrial pathway of apoptosis in human lymphoblasts and myeloma cells. *Carcinogenesis* 2005; 26(5): 883-91.
- [15] Muck AO, Seeger H, Wallwiener D. Class-specific pro-apoptotic effect of statins on human vascular endothelial cell. *Z Kardiol* 2004; 93: 398-402.
- [16] Castor CW, Bignell MC, Hossler PA, Roberts D. Connective tissue activation. XXI. Regulation of glycosaminoglycan metabolism by lymphocytes (CTAP-I) and platelet (CTAP-III) growth factors. *In Vitro Cell Dev Biol* 1981; 17: 777-85.
- [17] Castor CW, Prince RK, Dorstewitz EL. Characteristics of human fibroblasts cultivated *in vitro* from different anatomical sites. *Lab Invest* 1962; 11: 703-13.
- [18] Yaron I, Shirazi E, Yudovich R, Caspi D, Levartovsky D, Yaron M. Fluoxetine and amitriptyline inhibit nitric oxide, prostaglandin E2 and hyaluronic acid production in human synovial cells and synovial tissue cultures. *Arthritis Rheum* 1999; 42: 2561-8.
- [19] Roehm NW, Rodgers GH, Hatfield SM, Glasebrook AL. An improved colorimetric assay for cell proliferation and viability using the tetrazolium salt XTT. *J Immunol Methods* 1991; 142: 257-65.
- [20] Yokota K, Miyoshi F, Miyazaki T, et al. High concentration simvastatin induces apoptosis in fibroblast-like synoviocytes from patients with rheumatoid arthritis. *J Rheumatol* 2008; 35: 193-200.
- [21] Yim HW, Jong HS, Choi HH, et al. Cyclooxygenase-2 inhibits novel ginseng metabolite - mediated apoptosis. *Cancer Res* 2005; 65: 1952-60.
- [22] Pawlik A, Baskiewicz-Masiuk M, Machalinski B, Gawronska-Szklarz B. The effect of methotrexate and glucocorticosteroids on apoptosis of phythemaglutinin-stimulated mononuclear cells from peripheral blood. *Fundam Clin Pharmacol* 2005; 19: 81-5.
- [23] Catrina AI, Trollmo C, af Klint E, et al. Evidence that anti-tumor necrosis factor therapy with both etanercept and infliximab induces apoptosis in macrophages, but not lymphocytes, in rheumatoid arthritis joints: extended report. *Arthritis Rheum* 2005; 52: 61-72.
- [24] Rubins JB, Greatens T, Kratzke RA, Tan AT, Polunovsky VA, Bitterman P. Lovastatin induces apoptosis in malignant mesothelioma cells. *Am J Respir Crit Care Med* 1998; 157: 1616-22.
- [25] Jones KD, Couldwell WT, Hinton DR, et al. Lovastatin induces growth inhibition and apoptosis in human malignant glioma cells. *Biochem Biophys Res Commun* 1994; 205: 1681-7.
- [26] Vitale M, Di Matola T, Rossi G, Laezza C, Fenzi G, Bifulco M. Prenyltransferase inhibitors induce apoptosis in proliferating thyroid cells through a p53-independent, CrmA-sensitive, and caspase-3 like protease dependent mechanism. *Endocrinology* 1999; 140: 698-704.
- [27] Wong WWL, Tan MM, Xia Z, Dimitroulakos J, Minden MD, Penn LZ. Cerivastatin triggers tumor-specific apoptosis with higher efficacy than lovastatin. *Clin Cancer Res* 2001; 7: 2067-75.
- [28] Van de Donk NWCJ, Kamphuis MM, Lokhorst HM, Bloem AC. The cholesterol lowering drug lovastatin induces cell death in myeloma plasma cells. *Leukemia* 2002; 16: 1362-71.
- [29] Tan A, Levrey H, Dahm C, Polunovsky VA, Rubins J, Bitterman PB. Lovastatin induces fibroblast apoptosis *in vitro* and *in vivo*. A possible therapy for fibroproliferative disorders. *Am J Respir Crit Care Med* 1999; 159: 220-27.
- [30] Cafforio P, Dammacco F, Gernone A, Silvestris F. Statins activate the mitochondrial pathway of apoptosis in human lymphoblasts and myeloma cells. *Carcinogenesis* 2005; 26: 883-91.
- [31] Erl W, Hristov M, Neureuter M, Yan ZQ, Hansson GK, Weber PC. HMG-CoA reductase inhibitors induce apoptosis in neointima-derived vascular smooth muscle cells. *Atherosclerosis* 2003; 169(2): 251-8.
- [32] Nagashima T, Okazaki H, Yudoh K, Matsuno H, Minota S. Apoptosis of rheumatoid synovial cells by statins through the blocking of protein geranylgeranylation. *Arthritis Rheum* 2006; 54: 579-86.
- [33] Youssef S, Stuve O, Patarroyo JC, et al. The HMG-CoA reductase inhibitor, atorvastatin, promotes a Th2 bias and reverses paralysis in central nervous system autoimmune disease. *Nature* 2002; 420: 78-84.
- [34] Leung BP, Sattar N, Crilly A, et al. A novel anti-inflammatory role for simvastatin in inflammatory arthritis. *J Immunol* 2003; 170(3): 1524-30.
- [35] Del Rinco'n I, Williams K, Stern MP, Freeman GL, Escalante A. High incidence of cardiovascular events in a rheumatoid arthritis cohort not explained by traditional cardiac risk factors. *Arthritis Rheum* 2001; 44: 2737-45.