

Cellular Senescence is a Common Characteristic Shared by Preneoplastic and Osteo-Arthritic Tissue

Jean-Marc Brondello^{*1}, Didier Philipot¹, Farida Djouad¹, Christian Jorgensen^{1,2,3} and Danièle Noël¹

¹Inserm, U844, Montpellier, F-34295, France

²Université MONTPELLIER 1, UFR de Médecine, Montpellier, F-34000, France

³Immuno-Rheumatology Department, Lapeyronie Hospital, Montpellier, F-34295, France

Abstract: *Objective:* This study aims at highlighting the common signature between cartilaginous tissue in osteoarthritis (OA) and preneoplastic tissues preceding neoplasia and tumour formation and, second, focusing on the molecular mechanisms at the aetiology of both pathologies.

Results: Because age is the highest risk factor common for both OA and cancer development, it is tempting to compare the molecular mechanisms occurring at the onset of OA and preneoplastic lesions. Indeed, cellular senescence seems to be a common characteristic. Cellular senescence represents a natural barrier to suppress the unscheduled proliferation of damaged cells acting as a strong tumour suppressor pathway and in OA, it also occurs prematurely in chondrocytes. In this study, we review a number of molecular factors associated with the senescent phenotype.

Conclusion: Whereas accumulation of senescent cells in preneoplastic-like lesions leads to tissue degeneration and potentially tumour development; in OA, senescent cells accumulate in a slowly proliferative tissue. This is likely contributing at reducing the risk of cell transformation.

Keywords: Senescence, aging, chondrocytes, osteoarthritis, neoplasia.

INTRODUCTION

Osteoarthritis (OA) is a chronic degenerative disease characterized by progressive cartilage erosion leading to pain. At present, there is no efficient treatment to cure or even delay the progression of this most common disability-associated disease. The aetiology of OA is still unknown although clinical studies have shown that numerous factors are strong determinants. Among them, can be included genetic predisposition, obesity and primarily, age [1]. Remarkably, all of these factors are also associated with the prevalence of other age-related human diseases such as atherosclerosis [2] and preneoplastic lesions [3]. Recent works have focussed on the molecular events that govern the genesis of these lesions accumulating with age in numerous tissues [4, 5]. Preneoplastic lesions are benign. Nevertheless these cells may acquire, with time, new properties, such as genetic instability inducing *in fine* cellular transformation leading to neoplasia and eventually tumour formation. In summary, preneoplastic lesions are seen as natural barriers preventing tumour expansion and malignancy. Analysis of these recent studies reveals the strikingly high number of common characteristics between preneoplastic tissues and OA cartilage. Even if these common characteristics of cellular senescence in both pathologies may be merely co-incident, they have to be highlighted in the prospect of

using new therapeutic intervention in OA. Thus, the aim of this review is first, to highlight their common signatures and second, to focus on the molecular mechanisms shared by both pathologies.

ARTICULAR CARTILAGE BIOLOGY

The role of articular cartilage is to transmit forces across diarthrodial joints and maintain a friction-free surface at the epiphyseal ends of long bones to support limb movement. Articular cartilage is organised in various layers and contains one cell component, the chondrocyte which contributes by its anabolic function to the production and/or maintenance of the extracellular matrix (ECM) of the tissue [6]. The ECM is composed of a network of collagen fibres, primarily type II collagen and secondarily, type IX and XI collagens that give the tissue its shape, strength and tensile force. It also contains a highly hydrated gel of large proteoglycans, aggrecan being the most prominent, and glycoproteins which give resistance to compression [7]. Once the cartilage is formed in the adult, the turn-over of ECM components is very low.

OA PHYSIOPATHOLOGY

Among the numerous diseases which affect cartilage, OA is the most prevalent one. A number of studies have described the progression of this common disease [8]. The first detectable changes during OA by clinical examination are softening, fibrillation and ulceration of the cartilage. At latest stages of the disease, appearance of chondrocytes from neodifferentiated periosteal cells which eventually turn after calcification into osteophytes are present on the joint

*Address correspondence to this author at the Inserm U844, Hôpital Saint Eloi, Bâtiment INM, 80 Avenue Augustin Fliche, Montpellier, F-34295, France; Tel: 33 (0) 4 99 63 60 92; Fax: 33 (0) 4 99 63 60 20; E-mail: jean-marc.brondello@inserm.fr

edge [9]. These osteophytes, as a sort of extension of the articular surface, are responsible of the increasing pain. Finally, subchondral bone remodelling is proposed to be one driving force that accelerates cartilage [10].

Chondrocytes are the central players in the changes that occur during primary or secondary OA [8]. Anabolic function of chondrocytes is decreasing following pathology progression. This loss of function is mainly associated with a premature cellular senescence of chondrocytes [11]. Indeed, OA chondrocytes express a panel of senescence markers including acidic β -galactosidase activity [12], increase in reactive oxygen species (ROS), telomere attrition [13] and p53/p16^{ink4} accumulation [11]. Although the subject of debate, apoptotic chondrocytes have also been detected by tunnel assay in OA cartilage [14, 15]. Furthermore, an increased number of chondrocytes with altered phenotype and chondrocyte progenitors have also been found in OA cartilage [8, 16]. Finally, an increased number of hypertrophic chondrocytes, characterised by the expression of specific markers, such as collagen type X, alkaline phosphatase or runx2, is frequently observed [17]. In summary, OA is a multi-phenotypic complex disease where articular chondrocytes exhibit characteristics of stress-induced senescence with altered phenotypes and impaired normal anabolic capacities [11].

NEOPLASIA PHYSIOPATHOLOGY

Preneoplastic and neoplastic lesions are common. Both types of lesions are located within a high proliferative tissue and they increase with age. They are the result of abnormal cell proliferation that exceeds the surrounding normal tissue. Physiologically, preneoplasia precede neoplasia and both stages of lesion progression are hardly distinguishable. Neoplasia can remain benign or degenerate into tumour. Preneoplasia is induced *in vivo* following viral infection, recurrent genotoxic insults such as ultra-violet (UV), metabolic oxidative stress or genetic predisposition [18]. Genetic and molecular events that initiate this dysplasia have been broadly documented for the last 15 years. Indeed, these abnormal cells harbour an apparent attempt to hyperproliferate but rather are kept confined by a permanent cell cycle arrest associated with a senescent phenotype and impaired secretor activities [5, 19-21].

SIMILARITIES BETWEEN OA- AND PRENEOPLASIA/NEOPLASIA-ASSOCIATED MOLECULAR FACTORS

Cells are naturally exposed to various types of stress that lead to genomic assault during lifespan. Thus, cell-intrinsic mechanisms have evolved to limit the expansion of genetically altered cells. Cellular senescence is one of the natural barriers developed by multicellular organisms to suppress the unscheduled proliferation of damaged cells [19, 22, 23]. Cellular senescence is therefore a strong tumour suppressor pathway that is activated in benign preneoplastic-like lesions [5, 19, 21]. Remarkably, cellular senescence also occurs prematurely in OA chondrocytes. It is therefore tempting to make analogies between OA and preneoplasia associated characteristics (Fig. 1).

ROS and Associated Cellular Senescence

Oxidative damage and the chronic production of endogenous ROS have been associated with aging in various

human tissues and in animal models [24]. Articular chondrocytes actively produce ROS under mechanical stress or after inflammatory treatment [25, 26] and ROS secretion is believed to be one major mediator of OA-associated physiopathology [27]. Indeed, *in vitro* treatment of chondrocytes with oxygen peroxide (H₂O₂) used as source of ROS, resulted in p53/p21^{Waf1/Cip1}, p16^{ink4a} accumulation and telomere attrition which are both hallmarks of cellular senescence [12, 28, 29]. Furthermore, it has been reported that high level of ROS are produced by chondrocytes from OA patients [27, 30]. These molecular markers were also found in cartilage from patients with early OA suggesting their roles in the pathogenesis of the disease [11].

Interestingly, ROS have been proposed to play a pivotal role in the genesis of preneoplastic tissues [4]. Indeed, treatment of cells with ROS or transduction of primary epithelial cells or fibroblasts with oncogenes inducing preneoplasia *in vivo* led to the rapid increase of intracellular ROS production [31]. ROS participated to the DNA damage accumulation and thereby to the activation of the DNA damage response (DDR) [4]. The DDR is the signalling pathway orchestrated by ATM/ATR kinases (Ataxia telangiectasia mutated and ATM-Rad3 related) transducing toward two checkpoint kinases namely Chk1/Chk2. The DDR converges to p53 and induces senescence-dependent cell cycle arrest characteristic of preneoplastic cells. Therefore, DDR activation induced by ROS accumulation is now believed to be the hallmark of *in vivo* senescent tissues [22, 32].

Inflammation and Pro-Inflammatory Cytokines

Although OA is classified as a non inflammatory disease, pro-inflammatory cytokines such as IL-1 β , TNF- α or IL-8, have been reported to participate in OA physiopathology. Indeed, IL-1 β and IL-8 were detected both in the synovial fluid and in cartilage from OA patients [33, 34]. Depending on cell culture conditions, IL-1 β -treated chondrocytes underwent either dedifferentiation or telomere dependent-cellular senescence [29]. IL-8 has been proposed to induce chondrocyte hypertrophy and osteophyte formation which are observed in OA cartilage. Indeed, a time and dose-dependent increase in the expression level of specific hypertrophic markers was observed in OA chondrocytes treated with IL-8 [35]. Nevertheless, the role of this cytokine remains unclear.

For decades, IL-1 β and IL-8 are also associated with pre-malignant and malignant tumour lesions [36]. These inflammatory cytokines together with IL-6 were recently shown to play a crucial role in the premature cellular senescence found in preneoplastic lesions [36-38]. Indeed, when over-expressed in primary fibroblasts, IL-8 and one of its known receptor, CXCR2 were sufficient to induce premature senescence (acidic β -galactosidase staining and DDR activation) [38]. Furthermore, IL-8/CXCR2 tandem is transcriptionally induced by the DDR and participates to the senescence-dependent cell cycle arrest by producing more ROS [39]. Therefore, these signalling pathways including IL-8, IL-6 or IL-1 β were postulated to be tumour suppressors with pro-aging properties and accordingly found activated in benign lesions and inactivated during tumour progression [19, 38, 40, 41].




Fig. (1). Cellular senescence is a common characteristic of preneoplastic tissue and cartilage in early osteoarthritis (OA). **(A)** Neoplasia genesis. Upon environmental stress (carcinogens, oncogenes, genotoxics), in highly proliferative tissues such as epithelium, some cells will respond by abortive proliferation and emergence of senescent cells. If one cell overcomes the senescent process by inactivation of the DNA damage response (DDR), cell proliferation will lead to neoplasia and eventually tumour. **(B)** OA genesis. Upon environmental stress (mechanical stress, aging factors), OA chondrocytes become senescent and display an imbalance between anabolic and catabolic activity in favour of catabolism, leading to cartilage degradation. In both tissues, common senescence-associated markers are expressed: activated DDR and secretion of reactive oxygen species (ROS), interleukin (IL)-1 β , IL-8, insulin growth factor binding protein (IGFBP)-3, vascular endothelial growth factor (VEGF) and metalloproteinases (MMPs).

IGFBP Family Members

Differential analysis of gene and protein expression in OA versus healthy age-matched chondrocytes has revealed the over-expression of members of the IGFBP family such as insulin growth factor binding protein (IGFBP)-3 and -7 [42, 43]. IGFBP family members are pro-aging secreted factors known to modulate the insulin signalling pathway (for review, see [20]). Indeed, insulin pathway is central to aging and senescent mechanisms in modulating ROS production and cell survival [20, 44]. IGFBP-2/3 and -7 have also been shown to accumulate in preneoplastic-type lesions contributing to cellular senescence of these cells [3, 19, 45].

Vascularisation and VEGF Production

Another characteristic of OA cartilage is an increased expression level of vascular epithelial growth factor (VEGF). This growth factor is secreted by hypertrophic chondrocytes

and contributes to vascularisation and attraction of cartilage-resorbing cells during endochondral ossification in long bone development. In OA patients, VEGF production was shown to participate in the formation of osteophytes at the edges of articular cartilage [46]. However, the signalling pathways which control transcriptional induction of VEGF in OA chondrocytes are still being dissected. On the contrary, the transcriptional regulation and role of VEGF were broadly documented in early tumour development. Indeed, fibroblasts genetically modified to express oncogenes such as v-Ras or senescent fibroblasts were shown to secrete high levels of VEGF that can stimulate the proliferation of premalignant epithelial cells in culture and facilitate their conversion to malignant tumours *in vivo* [47]. This finding is believed to control the neo-vascularisation of neoplastic-like tissues and tumour cell survival to hypoxia-induced apoptosis.

Secretion of ECM Proteinases

Characteristic changes in early OA are similar to those seen with aging in cartilage from older, healthy subjects, with collagen damage and collagenases concentrated closer to the articular surface. Indeed, ECM degradation in OA was characterized by the secretion of metalloproteinases and aggrecanases such as MMP-1, -3, -8, -13 and a disintegrin and metalloproteinase with thrombospondin motif (ADAMTS)-5, respectively [48, 49]. Genetic inactivation of ADAMTS-5 prevented cartilage degeneration in OA-induced mice models [50, 51]. Recent studies revealed that senescent cells from preneoplastic tissues also promoted ECM degradation *via* the secretion of metalloproteinases such as MMP-3 [52]. The presence of metalloproteinases within preneoplastic lesions is believed to create a specific substratum which thereby facilitates the potential changes in cellular morphology required for tumour progression and metastasis [3].

CONCLUSIONS AND PERSPECTIVES

Cellular senescence seems to be a common process developed by OA chondrocytes and pre-tumour cells that may be at the genesis of both pathologies. Prevalence of these diseases increases with age and among the senescence-associated markers, ROS accumulation plays a central role. ROS production induces DDR-dependent premature senescence in preneoplastic cells and in OA chondrocytes contributing to the loss of their anabolic functions.

A number of studies now suggest that senescent cells secrete specific factors that set the stage for a cross-talk between senescent cells and their environment [20]. Such signals would be integrated into a coordinated response that can cause other cell components to prematurely senesce thereby establishing a pro-aging loop [20]. The consequence of the pro-aging loop is the degradation of the ECM, degeneration of the targeted tissue and production of survival factors such as VEGF [37]. Indeed, OA chondrocytes and preneoplastic-associated senescent cells both secrete VEGF but in pre-tumour tissues, VEGF helps tissue growth and survival by promoting new blood vessel formation [47]. In contrast, in the cartilage, the increase in vascularisation participates to the aberrant ossification of cartilage leading to osteophyte formation [46]. This phenomenon would favour subchondral bone pressure in OA patients.

Two exciting points of view emerge from all the similarities reported here between both diseases. First, as we mentioned, the aetiology seems to be similar. In proliferative tissue such as epithelium, the accumulation of senescent cells within preneoplastic-like lesions harbouring specific secretory properties will lead to tissue degeneration and may induce tumour development [5, 20]. By chance, accumulation of premature senescent chondrocytes in OA lesions occurs in a slowly proliferative tissue, dramatically reducing the risk of cell transformation yet promoting tissue degeneration. Second, research on OA physiopathology should take into account the literature on the deciphering of the neoplasia-associated molecular mechanisms and the prevention of tumour formation. Indeed, according to the numerous similarities between preneoplastic and OA lesions, it would be questionable whether the battery of pharmaceutical compounds available for anti-cancer

treatments such as anti-VEGF treatments could be tested by clinicians as potential OA treatments in order to delay or prevent the progression of this disease.

ACKNOWLEDGEMENTS

Work in the laboratory Inserm U844 is supported by the Inserm Institute, the University of Montpellier I and grants from the "Fondation Pour la Recherche Médicale".

REFERENCES

- [1] Felson DT, Lawrence RC, Dieppe PA, *et al.* Osteoarthritis: new insights. Part I: The disease and its risk factors. *Ann Intern Med* 2000; 133: 635-46.
- [2] Edo MD, Andres V. Aging, telomeres, and atherosclerosis. *Cardiovasc Res* 2005; 66: 213-21.
- [3] Campisi J, d'Adda di Fagnana F. Cellular senescence: When bad things happen to good cells? *Nat Rev Mol Cell Biol* 2007; 8: 729-40.
- [4] Bartek J, Bartkova J, Lukas J. DNA damage signalling guards against activated oncogenes and tumour progression. *Oncogene* 2007; 26: 7773-9.
- [5] Finkel T, Serrano M, Blasco MA. The common biology of cancer and ageing. *Nature* 2007; 448: 767-74.
- [6] Lefebvre V, Smits P. Transcriptional control of chondrocyte fate and differentiation. *Birth Defects Res C Embryo Today* 2005; 75: 200-12.
- [7] Poole AR, Kojima T, Yasuda T, Mwale F, Kobayashi M, Laverty S. Composition and structure of articular cartilage: a template for tissue repair. *Clin Orthop Relat Res* 2001; 391(Suppl): S26-33.
- [8] Aigner T, Soder S, Gebhard PM, McAlinden A, Haag J. Mechanisms of disease: Role of chondrocytes in the pathogenesis of osteoarthritis--structure, chaos and senescence. *Nat Clin Pract Rheumatol* 2007; 3: 391-9.
- [9] Buckwalter JA, Martin JA. Osteoarthritis. *Adv Drug Deliv Rev* 2006; 58: 150-67.
- [10] Martel-Pelletier J, Boileau C, Pelletier JP, Roughley PJ. Cartilage in normal and osteoarthritis conditions. *Best Pract Res Clin Rheumatol* 2008; 22: 351-84.
- [11] Loeser RF. Aging and osteoarthritis: the role of chondrocyte senescence and aging changes in the cartilage matrix. *Osteoarthritis Cartilage* 2009; 17: 971-79.
- [12] Martin JA, Buckwalter JA. The role of chondrocyte senescence in the pathogenesis of osteoarthritis and in limiting cartilage repair. *J Bone Joint Surg Am* 2003; 85-A(Suppl 2): 106-10.
- [13] Martin JA, Buckwalter JA. Telomere erosion and senescence in human articular cartilage chondrocytes. *J Gerontol A Biol Sci Med Sci* 2001; 56: B172-9.
- [14] Kim HA, Lee YJ, Seong SC, Choe KW, Song YW. Apoptotic chondrocyte death in human osteoarthritis. *J Rheumatol* 2000; 27: 455-62.
- [15] Mistry D, Oue Y, Chambers MG, Kayser MV, Mason RM. Chondrocyte death during murine osteoarthritis. *Osteoarthritis Cartilage* 2004; 12: 131-41.
- [16] Alsalameh S, Amin R, Gemba T, Lotz M. Identification of mesenchymal progenitor cells in normal and osteoarthritic human articular cartilage. *Arthritis Rheum* 2004; 50: 1522-32.
- [17] Girkontaite I, Frischholz S, Lammi P, *et al.* Immunolocalization of type X collagen in normal fetal and adult osteoarthritic cartilage with monoclonal antibodies. *Matrix Biol* 1996; 15: 231-8.
- [18] Besaratinia A, Pfeifer GP. Sunlight ultraviolet irradiation and BRAF V600 mutagenesis in human melanoma. *Hum Mutat* 2008; 29: 983-91.
- [19] Cichowski K, Hahn WC. Unexpected pieces to the senescence puzzle. *Cell* 2008; 133: 958-61.
- [20] Kuilman T, Peeper DS. Senescence-messaging secretome: SMS-ing cellular stress. *Nat Rev Cancer* 2009; 9: 81-94.
- [21] Mooi WJ, Peeper DS. Oncogene-induced cell senescence--halting on the road to cancer. *N Engl J Med* 2006; 355: 1037-46.
- [22] Di Micco R, Fumagalli M, Cicalese A, *et al.* Oncogene-induced senescence is a DNA damage response triggered by DNA hyper-replication. *Nature* 2006; 444: 638-42.
- [23] Prieur A, Peeper DS. Cellular senescence *in vivo*: A barrier to tumorigenesis. *Curr Opin Cell Biol* 2008; 20: 150-5.

- [24] Blagosklonny MV. Aging: ROS or TOR. *Cell Cycle* 2008; 7: 3344-54.
- [25] Chen AF, Davies CM, De Lin M, Fermor B. Oxidative DNA damage in osteoarthritic porcine articular cartilage. *J Cell Physiol* 2008; 217: 828-33.
- [26] Davies CM, Guilak F, Weinberg JB, Fermor B. Reactive nitrogen and oxygen species in interleukin-1-mediated DNA damage associated with osteoarthritis. *Osteoarthritis Cartilage* 2008; 16: 624-30.
- [27] Martin JA, Klingelhutz AJ, Moussavi-Harami F, Buckwalter JA. Effects of oxidative damage and telomerase activity on human articular cartilage chondrocyte senescence. *J Gerontol A Biol Sci Med Sci* 2004; 59: 324-37.
- [28] Zhou HW, Lou SQ, Zhang K. Recovery of function in osteoarthritic chondrocytes induced by p16INK4a-specific siRNA *in vitro*. *Rheumatology (Oxford)* 2004; 43: 555-68.
- [29] Dai SM, Shan ZZ, Nakamura H, et al. Catabolic stress induces features of chondrocyte senescence through overexpression of caveolin 1: Possible involvement of caveolin 1-induced down-regulation of articular chondrocytes in the pathogenesis of osteoarthritis. *Arthritis Rheum* 2006; 54: 818-31.
- [30] Yudoh K, Nguyen T, Nakamura H, Hongo-Masuko K, Kato T, Nishioka K. Potential involvement of oxidative stress in cartilage senescence and development of osteoarthritis: oxidative stress induces chondrocyte telomere instability and downregulation of chondrocyte function. *Arthritis Res Ther* 2005; 7: R380-91.
- [31] Collado M, Gil J, Efeyan A, et al. Tumour biology: senescence in premalignant tumours. *Nature* 2005; 436: 642.
- [32] Braig M, Lee S, Loddenkemper C, et al. Oncogene-induced senescence as an initial barrier in lymphoma development. *Nature* 2005; 436: 660-5.
- [33] Loeser RF. Molecular mechanisms of cartilage destruction: Mechanics, inflammatory mediators, and aging collide. *Arthritis Rheum* 2006; 54: 1357-60.
- [34] Fan Z, Bau B, Yang H, Soeder S, Aigner T. Freshly isolated osteoarthritic chondrocytes are catabolically more active than normal chondrocytes, but less responsive to catabolic stimulation with interleukin-1beta. *Arthritis Rheum* 2005; 52: 136-43.
- [35] Merz D, Liu R, Johnson K, Terkeltaub R. IL-8/CXCL8 and growth-related oncogene alpha/CXCL1 induce chondrocyte hypertrophic differentiation. *J Immunol* 2003; 171: 4406-15.
- [36] Kuilman T, Michaloglou C, Vredeveld LC, et al. Oncogene-induced senescence relayed by an interleukin-dependent inflammatory network. *Cell* 2008; 133: 1019-31.
- [37] Coppe JP, Patil CK, Rodier F, et al. Senescence-associated secretory phenotypes reveal cell-nonautonomous functions of oncogenic RAS and the p53 tumor suppressor. *PLoS Biol* 2008; 6: 2853-68.
- [38] Acosta JC, O'Loughlen A, Banito A, et al. Chemokine signaling via the CXCR2 receptor reinforces senescence. *Cell* 2008; 133: 1006-18.
- [39] Acosta JC, O'Loughlen A, Banito A, Raguz S, Gil J. Control of senescence by CXCR2 and its ligands. *Cell Cycle* 2008; 7: 2956-9.
- [40] Acosta JC, Gil J. A role for CXCR2 in senescence, but what about in cancer? *Cancer Res* 2009; 69: 2167-70.
- [41] Rodier F, Coppe JP, Patil CK, et al. Persistent DNA damage signalling triggers senescence-associated inflammatory cytokine secretion. *Nat Cell Biol* 2009; 11: 973-9.
- [42] Aigner T, Fundel K, Saas J, et al. Large-scale gene expression profiling reveals major pathogenetic pathways of cartilage degeneration in osteoarthritis. *Arthritis Rheum* 2006; 54: 3533-44.
- [43] Iliopoulos D, Malizos KN, Oikonomou P, Tsezou A. Integrative microRNA and proteomic approaches identify novel osteoarthritis genes and their collaborative metabolic and inflammatory networks. *PLoS ONE* 2008; 3: e3740.
- [44] Greer EL, Brunet A. Signaling networks in aging. *J Cell Sci* 2008; 121: 407-12.
- [45] Wajapeyee N, Serra RW, Zhu X, Mahalingam M, Green MR. Oncogenic BRAF induces senescence and apoptosis through pathways mediated by the secreted protein IGFBP7. *Cell* 2008; 132: 363-74.
- [46] Murata M, Yudoh K, Masuko K. The potential role of vascular endothelial growth factor (VEGF) in cartilage: How the angiogenic factor could be involved in the pathogenesis of osteoarthritis? *Osteoarthritis Cartilage* 2008; 16: 279-86.
- [47] Coppe JP, Kauer K, Campisi J, Beausejour CM. Secretion of vascular endothelial growth factor by primary human fibroblasts at senescence. *J Biol Chem* 2006; 281: 29568-74.
- [48] Hollander AP, Pidoux I, Reiner A, Rorabeck C, Bourne R, Poole AR. Damage to type II collagen in aging and osteoarthritis starts at the articular surface, originates around chondrocytes, and extends into the cartilage with progressive degeneration. *J Clin Invest* 1995; 96: 2859-69.
- [49] Wu W, Billingham RC, Pidoux I, et al. Sites of collagenase cleavage and denaturation of type II collagen in aging and osteoarthritic articular cartilage and their relationship to the distribution of matrix metalloproteinase 1 and matrix metalloproteinase 13. *Arthritis Rheum* 2002; 46: 2087-94.
- [50] Glasson SS, Askew R, Sheppard B, et al. Deletion of active ADAMTS5 prevents cartilage degradation in a murine model of osteoarthritis. *Nature* 2005; 434: 644-8.
- [51] Stanton H, Rogerson FM, East CJ, et al. ADAMTS5 is the major aggrecanase in mouse cartilage *in vivo* and *in vitro*. *Nature* 2005; 434: 648-52.
- [52] Parrinello S, Coppe JP, Krtolica A, Campisi J. Stromal-epithelial interactions in aging and cancer: Senescent fibroblasts alter epithelial cell differentiation. *J Cell Sci* 2005; 118: 485-96.

Received: December 6, 2009

Revised: January 6, 2010

Accepted: January 18, 2010

© Brondello et al.; licensee Bentham Open.

This is an open access article licensed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/3.0/>) which permits unrestricted, non-commercial use, distribution and reproduction in any medium, provided the work is properly cited.